

For this assignment, you should copy this document as closely as possible. Put your name in the top right corner. These are some concepts from Chapter 0 that you should already know.

- Definition (*The Well Ordering Principle*)** - Every nonempty set of positive integers contains a smallest member.
- Theorem (*The Division Algorithm*)** - Let a and b be integers with $b > 0$. Then there exist unique integers q and r with the property that $a = bq + r$, where $0 \leq r < b$.
- Definition** - The **Greatest Common Divisor** of two nonzero integers a and b is the largest of all common divisors of a and b . We denote this integer by $\gcd(a, b)$. When $\gcd(a, b) = 1$, we say a and b are *relatively prime*.
- Theorem** For any nonzero integers a and b , there exist integers s and t such that $\gcd(a, b) = as + bt$. Moreover, $\gcd(a, b)$ is the smallest positive integer of the form $as + bt$.
- Corollary** If a and b are relatively prime, then there exist integers s and t such that $as + bt = 1$.
- Theorem (*Euclid's Lemma*)** If p is a prime that divides ab , then p divides a or p divides b (or both).
Proof: Suppose that p is a prime that divides ab , but without loss of generality (WLOG) does not divide a . Then we must show that p divides b . Since p does not divide a , then a and p are relatively prime. So there exist integers s and t such that $1 = as + pt$. Multiply through by b to get $b = abs + ptb$. Since p divides ab and p divides a , p divides the right hand side of the equation. Hence p divides the left as well. So p divides b . \square .
- Theorem (*Fundamental Theorem of Arithmetic*)** Every integer greater than 1 is a prime or a product of primes. This product is unique, except for the order in which the factors appear. That is, if $n = p_1 p_2 \dots p_r$ and $n = q_1 q_2 \dots q_s$, where the p 's and q 's are primes, then $r = s$ and, after renumbering the q 's, we have $p_i = q_i$ for all i .
- Definition** The *least common multiple* of two nonzero integers a and b is the smallest positive integer that is a multiple of both a and b . We denote this integer by $\text{lcm}(a, b)$.
- Theorem (*The First Principle of Mathematical Induction*)** Let S be a set of integers containing a . Suppose S has the property that whenever some integer $n \geq a$ belongs to S , then the integer $n + 1$ belongs to S . Then S contains every integer greater than or equal to a .
- Theorem (*DeMoivre's Theorem*)** For every positive integer n and every real number θ , $(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$, where i is the complex number $\sqrt{-1}$.
Proof: Base Step: The statement is clearly true for $n = 1$.
 Inductive Step: Assume true for $n = 1$. Show the statement is true for $n + 1$. In other words, assume $(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$, prove $(\cos \theta + i \sin \theta)^{(n+1)} = \cos(n+1)\theta + i \sin(n+1)\theta$. We see that

$$\begin{aligned} (\cos \theta + i \sin \theta)^n &= (\cos \theta + i \sin \theta)^{(n+1)} (\cos \theta + i \sin \theta) & (1) \\ &= (\cos n\theta + i \sin n\theta)(\cos \theta + i \sin \theta) & (2) \\ &= \cos n\theta \cos \theta + i(\sin n\theta \cos \theta + \sin \theta \cos n\theta) - \sin n\theta \sin \theta. & (3) \end{aligned}$$
 Now, using trig identities for $\cos(\alpha + \beta)$ and $\sin(\alpha + \beta)$, we see that this last term is $\cos(n+1)\theta + i \sin(n+1)\theta$. So, by induction, the statement is true for all positive integers. \square
- Theorem (*The Second (Strong) Principle of Mathematical Induction*)** Let S be a set of integers containing a . Suppose S has the property that n belongs to S whenever every integer less than n and greater than or equal to a belongs to S . Then S contains every integer greater than or equal to a .

12. **Definition** An *equivalence relation* on a set S is a set R of ordered pairs of elements of S such that
- (a) $(a, a) \in R$ for all $a \in S$. (reflexive property)
 - (b) $(a, b) \in R$ implies $(b, a) \in R$ (symmetric property)
 - (c) $(a, b) \in R$ and $(b, c) \in R$ imply $(a, c) \in R$ (transitive property)
13. **Definition** A *partition* of a set S is a collection of nonempty disjoint subsets of S whose union is S .
14. **Theorem** The equivalence classes of an equivalence relation on a set S constitute a partition of S . Conversely, for any partition P of S , there is an equivalence relation on S whose equivalence classes are the elements of P .

If these concepts do NOT look familiar. Be sure to study them carefully! Your homework is the following from Chapter 0: 2, 3, 4, 7, 8, 9, 11, 14, 20, 21, 22, 30, 34, 53, 54.