

MATH 231, Chapter 0

Calculus with Functions

James Madison University

Intervals - describe the set of real numbers between -2 and 4, including -2 but not including 4:



Intervals - describe the set of real numbers between -2 and 4, including -2 but not including 4:



$$\{x \mid -2 \leq x < 4\}$$

Intervals - describe the set of real numbers between -2 and 4, including -2 but not including 4:



$$\{x \mid -2 \leq x < 4\} = [-2, 4)$$

Intervals - describe the set of real numbers between -2 and 4, including -2 but not including 4:



$\{x \mid -2 \leq x < 4\} = [-2, 4)$
(the last, $[-2, 4)$, is *interval notation*)

Intervals - describe the set of real numbers between -2 and 4, including -2 but not including 4:



$\{x \mid -2 \leq x < 4\} = [-2, 4)$
(the last, $[-2, 4)$, is *interval notation*)

Sets?

Intervals - describe the set of real numbers between -2 and 4, including -2 but not including 4:



$\{x \mid -2 \leq x < 4\} = [-2, 4)$
(the last, $[-2, 4)$, is *interval notation*)

Sets?

$\{5, 11, -3\}$

Intervals - describe the set of real numbers between -2 and 4, including -2 but not including 4:



$\{x \mid -2 \leq x < 4\} = [-2, 4)$
(the last, $[-2, 4)$, is *interval notation*)

Sets?

$\{5, 11, -3\}$

$\{3, 6, 9, 12, \dots\}$

Intervals - describe the set of real numbers between -2 and 4, including -2 but not including 4:



$\{x \mid -2 \leq x < 4\} = [-2, 4)$
(the last, $[-2, 4)$, is *interval notation*)

Sets?

$\{5, 11, -3\}$

$\{3, 6, 9, 12, \dots\}$

Set Notation and Elements of a Set

Schematically, **set notation** for a set S is written in the form

$$S = \{x \in \text{category} \mid \text{test determining whether } x \text{ is in the set}\}.$$

An object x is an **element** of a set S if it is contained in S , that is, if it passes the test written in the second half of the preceding notation. When this happens, we write $x \in S$.

Write the following in interval notation:




Write the following in interval notation:



Union and Intersection

Suppose A and B are sets.

- (a) The **union** of A and B is the set $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.
- (b) The **intersection** of A and B is the set $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$.




Union and Intersection

Suppose A and B are sets.

(a) The **union** of A and B is the set $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.

(b) The **intersection** of A and B is the set $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$.




\in , \subset , \emptyset

Union and Intersection

Suppose A and B are sets.

(a) The **union** of A and B is the set $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$.

(b) The **intersection** of A and B is the set $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$.



\in , \subset , \emptyset



Rational and Irrational Numbers

(a) A rational number is a real number that can be written as a quotient of the form $\frac{p}{q}$ for some integers p and q , with $q \neq 0$.

(b) An irrational number is a real number that cannot be written in the form of a rational number.

For the record, some standard sets of real numbers will be designated as follows.

For the record, some standard sets of real numbers will be designated as follows.

- ▶ The set of natural numbers = $\{1, 2, 3, 4, \dots\}$ - \mathbb{N} .

For the record, some standard sets of real numbers will be designated as follows.

- ▶ The set of natural numbers = $\{1, 2, 3, 4, \dots\}$ - \mathbb{N} .
- ▶ The set of whole numbers = $\{0, 1, 2, 3, \dots\}$ - \mathbb{W} .

Review the basic language of sets and the meaning of symbols like \in , \subset , \cup , \cap , \emptyset , \sim and so on.

For the record, some standard sets of real numbers will be designated as follows.

- ▶ The set of natural numbers = $\{1, 2, 3, 4, \dots\}$ - \mathbb{N} .
- ▶ The set of whole numbers = $\{0, 1, 2, 3, \dots\}$ - \mathbb{W} .
- ▶ The set of integers = $\{0, \pm 1, \pm 2, \pm 3, \dots\}$ - \mathbb{Z}

Review the basic language of sets and the meaning of symbols like \in , \subset , \cup , \cap , \emptyset , \sim and so on.

For the record, some standard sets of real numbers will be designated as follows.

- ▶ The set of natural numbers = $\{1, 2, 3, 4, \dots\}$ - \mathbb{N} .
- ▶ The set of whole numbers = $\{0, 1, 2, 3, \dots\}$ - \mathbb{W} .
- ▶ The set of integers = $\{0, \pm 1, \pm 2, \pm 3, \dots\}$ - \mathbb{Z}
- ▶ The set of rational numbers = $\{x \mid x = \frac{p}{q}, p, q \in \mathbb{Z}, q \neq 0\}$ - \mathbb{Q}


Review the basic language of sets and the meaning of symbols like \in , \subset , \cup , \cap , \emptyset , \sim and so on.

For the record, some standard sets of real numbers will be designated as follows.

- ▶ The set of natural numbers = $\{1, 2, 3, 4, \dots\}$ - \mathbb{N} .
- ▶ The set of whole numbers = $\{0, 1, 2, 3, \dots\}$ - \mathbb{W} .
- ▶ The set of integers = $\{0, \pm 1, \pm 2, \pm 3, \dots\}$ - \mathbb{Z}
- ▶ The set of rational numbers = $\{x \mid x = \frac{p}{q}, p, q \in \mathbb{Z}, q \neq 0\}$ - \mathbb{Q}
- ▶ The set of real numbers - \mathbb{R} .

Review the basic language of sets and the meaning of symbols like \in , \subset , \cup , \cap , \emptyset , \sim and so on.

Cartesian co-ordinates



Distance between two points: $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

Write the equation of a circle with center $(2, -1)$ and radius 3.

Are you familiar with the other *conic sections*?

Solving (mostly polynomial) equations.

Solving (mostly polynomial) equations.

Solve:

▶ $2x - 3 = 9$

Solving (mostly polynomial) equations.

Solve:

▶ $2x - 3 = 9$

▶ $2x^2 - 5x + 3 = 0$

Solving (mostly polynomial) equations.

Solve:

▶ $2x - 3 = 9$

▶ $2x^2 - 5x + 3 = 0$

▶ $x^2 + x = 2$

Solving (mostly polynomial) equations.

Solve:

▶ $2x - 3 = 9$

▶ $2x^2 - 5x + 3 = 0$

▶ $x^2 + x = 2$

▶ $x^2 + x = 3$

Solving (mostly polynomial) equations.

Solve:

▶ $2x - 3 = 9$

▶ $2x^2 - 5x + 3 = 0$

▶ $x^2 + x = 2$

▶ $x^2 + x = 3$

▶ $\frac{2x^2 - 5x + 3}{x^2 - 2} = 0$

Solving (mostly polynomial) equations.

Solve:

▶ $2x - 3 = 9$

▶ $2x^2 - 5x + 3 = 0$

▶ $x^2 + x = 2$

▶ $x^2 + x = 3$

▶ $\frac{2x^2 - 5x + 3}{x^2 - 2} = 0$

▶ $\frac{2x^2 - 5x + 3}{x^2 - 1} = 0$

The Quadratic Formula

If a , b , and c are real numbers, the solutions of the quadratic equation $ax^2 + bx + c = 0$ are of the form

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Formulas for Factoring Differences of Powers

For all real numbers a and b , and any positive integer n ,

$$a^2 - b^2 = (a - b)(a + b)$$

$$a^3 - b^3 = (a - b)(a^2 + ab + b^2)$$

$$a^n - b^n = (a - b)(a^{n-1} + a^{n-2}b + a^{n-3}b^2 + a^{n-4}b^3 + \dots + a^2b^{n-3} + ab^{n-2} + b^{n-1})$$

Formulas for Factoring Differences of Powers

For all real numbers a and b , and any positive integer n ,

$$a^2 - b^2 = (a - b)(a + b)$$

$$a^3 - b^3 = (a - b)(a^2 + ab + b^2)$$

$$a^n - b^n = (a - b)(a^{n-1} + a^{n-2}b + a^{n-3}b^2 + a^{n-4}b^3 + \dots + a^2b^{n-3} + ab^{n-2} + b^{n-1})$$

Algebraic Rules for Fractions

Let a , b , c , and d be any real numbers. Then (assuming that no denominator in any expression is zero)

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd}$$

$$\left(\frac{a}{b}\right)\left(\frac{c}{d}\right) = \frac{ac}{bd}$$

$$\frac{(a/b)}{(c/d)} = \left(\frac{a}{b}\right)\left(\frac{d}{c}\right) = \frac{ad}{bc}$$

Formulas for Factoring Differences of Powers

For all real numbers a and b , and any positive integer n ,

$$a^2 - b^2 = (a - b)(a + b)$$

$$a^3 - b^3 = (a - b)(a^2 + ab + b^2)$$

$$a^n - b^n = (a - b)(a^{n-1} + a^{n-2}b + a^{n-3}b^2 + a^{n-4}b^3 + \dots + a^2b^{n-3} + ab^{n-2} + b^{n-1})$$

Algebraic Rules for Fractions

Let a , b , c , and d be any real numbers. Then (assuming that no denominator in any expression is zero)

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd} \qquad \left(\frac{a}{b}\right)\left(\frac{c}{d}\right) = \frac{ac}{bd} \qquad \frac{(a/b)}{(c/d)} = \left(\frac{a}{b}\right)\left(\frac{d}{c}\right) = \frac{ad}{bc}$$

More Algebraic Rules for Fractions

Let a , b , c , and d be any real numbers. Then (assuming that no denominator in any expression is zero)

$$\frac{a}{b} - \frac{c}{d} = \frac{ad - bc}{bd} \qquad c\left(\frac{a}{b}\right) = \frac{ac}{b} \qquad \frac{(a/b)}{c} = \frac{a}{bc} \qquad \frac{a}{(b/c)} = \frac{ac}{b}$$

Systems of equations?

$$\begin{cases} x - y = -1 \\ y + x^2 = 3 \end{cases}$$

Solving (mostly polynomial) inequalities.

Solving (mostly polynomial) inequalities.

Algebraic Rules for Inequalities

Suppose a , b , and c are nonzero real numbers.

- (a) If $a < b$ and $c > 0$, then $ac < bc$. (b) If $a < b$ and $c < 0$, then $ac > bc$.
- (c) If $0 < a < b$, then $\frac{1}{a} > \frac{1}{b}$. (d) If $a < b$, then $a + c < b + c$.

▶ $2x - 3 < 9$

▶ $x^2 + x \leq 2$

▶ $2x - 3 < 9$

▶ $x^2 + x \leq 2$

▶ $\frac{1}{x+3} \geq -2$

▶ $2x - 3 < 9$

▶ $x^2 + x \leq 2$

▶ $\frac{1}{x+3} \geq -2$

▶ $|x - 3| < 3$

▶ $|2x - 1| \geq 5$

Write the following in interval notation:

$$\{x ; x < 5\} =$$

Write the following in interval notation:

$$\{x ; x < 5\} =$$

$$\{x ; x \geq -3\} =$$

Write the following in interval notation:

$$\{x ; x < 5\} =$$

$$\{x ; x \geq -3\} =$$

$$\{x ; 3x - 5 \geq 0\} =$$

Write the following in interval notation:

$$\{x ; x < 5\} =$$

$$\{x ; x \geq -3\} =$$

$$\{x ; 3x - 5 \geq 0\} =$$

$$\{x ; 1 < -3x + 5 \leq 2\} =$$

Write the following in interval notation:

$$\{x ; x < 5\} =$$

$$\{x ; x \geq -3\} =$$

$$\{x ; 3x - 5 \geq 0\} =$$

$$\{x ; 1 < -3x + 5 \leq 2\} =$$

$$\{x ; |2x - 7| < 3\} =$$

Write the following in interval notation:

$$\{x ; x < 5\} =$$

$$\{x ; x \geq -3\} =$$

$$\{x ; 3x - 5 \geq 0\} =$$

$$\{x ; 1 < -3x + 5 \leq 2\} =$$

$$\{x ; |2x - 7| < 3\} =$$

$$\{x ; |2x - 7| > 3\} =$$

Functions

A **function** f from a set A to a set B is an assignment f that associates to each element x of the **domain** set A exactly one element $f(x)$ of the **codomain**, or **target**, set B .

Functions

A **function** f from a set A to a set B is an assignment f that associates to each element x of the **domain** set A exactly one element $f(x)$ of the **codomain**, or **target**, set B .

Domain and Range of a Function

If f is a function between unspecified subsets of \mathbb{R} , then we will take the **domain** of f to be the largest subset of \mathbb{R} for which f is defined:

$$\text{Domain}(f) = \{x \in \mathbb{R} \mid f(x) \text{ is defined}\}.$$

The **range** of such a function is the set of all possible outputs that it can attain:

$$\text{Range}(f) = \{y \in \mathbb{R} \mid \text{there is some } x \in \text{Domain}(f) \text{ for which } f(x) = y\}.$$

One-to-One Function

A function f is *one-to-one* if, for all a and b in the domain of f ,

$$a \neq b \implies f(a) \neq f(b).$$

One-to-One Function

A function f is *one-to-one* if, for all a and b in the domain of f ,


$$a \neq b \implies f(a) \neq f(b).$$

The Graph of a Function


The *graph* of a function f is the collection of ordered pairs $(x, f(x))$ for which x is in the domain of f . In set notation we can write

$$\text{Graph}(f) = \{ (x, f(x)) \mid x \in \text{Domain}(f) \}.$$


A graph that is a function



A graph that is not a function



A function, but not one-to-one



Example - let f be the function that assigns each real number to its square.


$$f : x \rightarrow x^2$$

Example - let f be the function that assigns each real number to its square.

$$f : x \rightarrow x^2$$

or

$$f(x) = x^2$$




Example - let f be the function that assigns each real number to its square.

$$f : x \rightarrow x^2$$

or

$$f(x) = x^2$$




f - "the function"

Example - let f be the function that assigns each real number to its square.

$$f : x \rightarrow x^2$$

or

$$f(x) = x^2$$



f - "the function"

$f(x)$ the number that the function assigns to some input value x .

$$f(x) = x^2$$

Domain of f ?

Range of f ?

$$f(x) = x^2$$

Domain of f ?

Range of f ?

Find

▶ $f(3)$

▶ $f(w - 2)$

▶ $\frac{f(z)}{f(w^3+1)+2}$

▶ $\frac{f(x+h)-f(x)}{h}$

Suppose:

$$\blacktriangleright g(x) = \sqrt{x^2 + 2x}$$

$$\blacktriangleright h(x) = \frac{x^2}{x^2+1}$$

$$\blacktriangleright k(x) = \frac{2x}{9x^2-4}$$

Suppose:

$$\blacktriangleright g(x) = \sqrt{x^2 + 2x}$$

$$\blacktriangleright h(x) = \frac{x^2}{x^2+1}$$

$$\blacktriangleright k(x) = \frac{2x}{9x^2-4}$$

Domain and Range for g , h , k ?

Suppose:

$$\blacktriangleright g(x) = \sqrt{x^2 + 2x}$$

Domain and Range for g ?

Suppose:



$$\blacktriangleright h(x) = \frac{x^2}{x^2+1}$$



Domain and Range for h ?

Suppose:



$$\blacktriangleright k(x) = \frac{2x}{9x^2 - 4}$$

Domain and Range for k ?





What are the domain and range of f ?




What are the domain and range of f ? g ?

Vocabulary	Definition	Behavior
f has a root at $x = c$	$f(c) = 0$	graph intersects the x -axis at $x = c$
f has a y-intercept at $y = b$	$f(0) = b$	graph intersects the y -axis at $y = b$
f is positive on I	$f(x) > 0$ for all $x \in I$	graph is above the x -axis on I
f is increasing on I	$f(b) > f(a)$ for all $b > a$ in I	graph moves up as we look from left to right on I
f has a local maximum at $x = c$	$f(c) \geq f(x)$ for all x near $x = c$	graph has a relative “hilltop” at $x = c$
f has a global maximum at $x = c$	$f(c) \geq f(x)$ for all $x \in \text{Domain}(f)$	graph is the highest at $x = c$
f is concave up on I	<i>will state precisely in Section 3.3</i>	graph curves upwards on I like part of a “U”
f has an inflection point at $x = c$	<i>will state precisely in Section 3.3</i>	graph of f changes concavity at $x = c$

Algebraic functions

Type	General Form	Examples
Linear	$f(x) = mx + b$, where m and b are any real numbers	$f(x) = 2x - 1$, $f(x) = 1.4x$
Power	$f(x) = Ax^k$, where $A \neq 0$ is real and k is rational	$f(x) = 3x^3$, $f(x) = 1.7x^{-1/5}$
Polynomial	$f(x) = a_nx^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$, where n is a nonnegative integer and each a_i is a real number	$f(x) = 3x^5 - 2x^3 + x - 6$
Rational	$f(x) = \frac{p(x)}{q(x)}$, where $p(x)$ and $q(x)$ are polynomials	$f(x) = \frac{3x^2 - 1}{x^5 + x^3 + 1}$
Other	any other algebraic function that is not one of the types listed	$f(x) = \frac{1 + \sqrt{x}}{1 + x}$, $f(x) = x^{2/3} + 5$



Straight lines

$$y = mx + b$$

$$m = \text{slope} = \text{"rate of change"} = \frac{\text{vertical change}}{\text{horizontal change}} = \frac{y_2 - y_1}{x_2 - x_1}$$

b = y -axis intercept

Point slope form $y - y_1 = m(x - x_1)$


“General” form $Ax + By + C = 0$

Horizontal line ?


Vertical line ?

Find the equation of each line graphed below:


a.




b.



c.



d.




Also find:

An equation for the line parallel to a) through $(2, 1)$.

An equation for the line perpendicular to b) through $(-3, 2)$.


a.



Also find:

An equation for the line parallel to a) through $(2, 1)$.


b.




Also find:

An equation for the line perpendicular to b) through $(-3, 2)$.

C.



d.



Transcendental functions

Type	Examples
Exponential	$f(x) = 2^x$, $f(x) = 3e^{4x}$, $f(x) = 1.2(3.4)^x$
Logarithmic	$f(x) = \log_{10} x$, $f(x) = \ln x$, $f(x) = \log_2 x$
Trigonometric	$f(x) = \sin x$, $f(x) = \cos x$, $f(x) = \cot x$
Inverse Trigonometric	$f(x) = \arcsin x$, $f(x) = \cos^{-1} x$, $f(x) = \arctan x$
Other	$f(x) = x + \sin x$, $f(x) = \ln(\sqrt{x} + 12)$, $f(x) = 2^x \arctan x$

Arithmetic Combinations of Functions

Suppose f and g are functions and k is a real number.

- (a) The **constant multiple** of f by k is the function kf defined by $(kf)(x) = kf(x)$ for all x in the domain of f .
- (b) The **sum** of f and g is the function $f + g$ defined by $(f + g)(x) = f(x) + g(x)$ for all x in the domains of both f and g .
- (c) The **product** of f and g is the function $f \cdot g$ defined by $(f \cdot g)(x) = f(x)g(x)$ for all x in the domains of both f and g .
- (d) The **quotient** of f and g is the function $\frac{f}{g}$ defined by $\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$ for all x in the domains of both f and g with $g(x) \neq 0$.

The Composition of Two Functions

The **composition** of two functions f and g is the function $f \circ g$ defined by

$$(f \circ g)(x) = f(g(x))$$

for all x in the domain of g such that $g(x)$ is in the domain of f .

Transformation	Graphical Result	Algebraic Result
$f(x) + C$	shifts up C units if $C > 0$ shifts down C units if $C < 0$	$(x, y) \rightarrow (x, y + C)$
$f(x + C)$	shifts left C units if $C > 0$ shifts right C units if $C < 0$	$(x, y) \rightarrow (x - C, y)$
$kf(x)$	vertical stretch by k if $k > 1$ vertical compression by k if $0 < k < 1$	$(x, y) \rightarrow (x, ky)$
$f(kx)$	horizontal compression by k if $k > 1$ horizontal stretch by k if $0 < k < 1$	$(x, y) \rightarrow \left(\frac{1}{k}x, y\right)$
$-f(x)$	graph reflects across the x -axis	$(x, y) \rightarrow (x, -y)$
$f(-x)$	graph reflects across the y -axis	$(x, y) \rightarrow (-x, y)$

Suppose:

▶ $f(x) = x^2$

▶ $g(x) = \sqrt{x^2 + 2x}$

▶ $h(x) = \frac{x^2}{x^2+1}$

▶ $k(x) = \frac{2x}{9x^2-4}$

Suppose:

▶ $f(x) = x^2$

▶ $g(x) = \sqrt{x^2 + 2x}$

▶ $h(x) = \frac{x^2}{x^2+1}$

▶ $k(x) = \frac{2x}{9x^2-4}$

Find

▶ $(f + g)(x)$

Suppose:

$$\blacktriangleright f(x) = x^2$$

$$\blacktriangleright g(x) = \sqrt{x^2 + 2x}$$

$$\blacktriangleright h(x) = \frac{x^2}{x^2+1}$$

$$\blacktriangleright k(x) = \frac{2x}{9x^2-4}$$

Find

$$\blacktriangleright (f + g)(x)$$

$$\blacktriangleright \frac{h}{g}(x) = \frac{h(x)}{g(x)}$$

Suppose:

$$\blacktriangleright f(x) = x^2$$

$$\blacktriangleright g(x) = \sqrt{x^2 + 2x}$$

$$\blacktriangleright h(x) = \frac{x^2}{x^2+1}$$

$$\blacktriangleright k(x) = \frac{2x}{9x^2-4}$$

Find

$$\blacktriangleright (f + g)(x)$$

$$\blacktriangleright \frac{h}{g}(x) = \frac{h(x)}{g(x)}$$

$$\blacktriangleright k \circ f(x) = k(f(x))$$

Suppose:

▶ $f(x) = x^2$

▶ $g(x) = \sqrt{x^2 + 2x}$

▶ $h(x) = \frac{x^2}{x^2+1}$

▶ $k(x) = \frac{2x}{9x^2-4}$

Find

▶ $(f + g)(x)$

▶ $\frac{h}{g}(x) = \frac{h(x)}{g(x)}$

▶ $k \circ f(x) = k(f(x))$

▶ $fg(x)$ (or $(f * g)(x)$)

Suppose:

$$\blacktriangleright f(x) = x^2$$

$$\blacktriangleright g(x) = \sqrt{x^2 + 2x}$$

$$\blacktriangleright h(x) = \frac{x^2}{x^2+1}$$

$$\blacktriangleright k(x) = \frac{2x}{9x^2-4}$$

Find

$$\blacktriangleright (f + g)(x)$$

$$\blacktriangleright \frac{h}{g}(x) = \frac{h(x)}{g(x)}$$

$$\blacktriangleright k \circ f(x) = k(f(x))$$

$$\blacktriangleright fg(x) \text{ (or } (f * g)(x))$$

$$\blacktriangleright g \circ f(x) = g(f(x))$$

Suppose:

$$\blacktriangleright f(x) = x^2$$

$$\blacktriangleright g(x) = \sqrt{x^2 + 2x}$$

$$\blacktriangleright h(x) = \frac{x^2}{x^2+1}$$

$$\blacktriangleright k(x) = \frac{2x}{9x^2-4}$$

Find

$$\blacktriangleright (f + g)(x)$$



$$\blacktriangleright \frac{h}{g}(x) = \frac{h(x)}{g(x)}$$



$$\blacktriangleright k \circ f(x) = k(f(x))$$

$$\blacktriangleright fg(x) \text{ (or } (f * g)(x))$$

$$\blacktriangleright g \circ f(x) = g(f(x))$$

$$\blacktriangleright f \circ g(x) = f(g(x))$$





Find

- ▶ $(f + g)(2)$
- ▶ $\frac{f}{g}(2)$
- ▶ $g \circ f(3)$
- ▶ $fg(2)$
- ▶ $f \circ g(2)$ (approx.)

Even and Odd Functions

A function f is an ***even function*** if $f(-x) = f(x)$ for all x in the domain of f .


A function f is an ***odd function*** if $f(-x) = -f(x)$ for all x in the domain of f .

Even and Odd Functions


A function f is an **even function** if $f(-x) = f(x)$ for all x in the domain of f .

A function f is an **odd function** if $f(-x) = -f(x)$ for all x in the domain of f .


$f(x) = \frac{1}{x}$
is an odd function



$g(x) = x^4 - x^2$
is an even function



$h(x) = \frac{2+x}{1+x^2}$
is neither even nor odd



The Inverse of a Function

If f and g are functions such that

$$g(f(x)) = x, \text{ for all } x \text{ in the domain of } f$$

$$f(g(x)) = x, \text{ for all } x \text{ in the domain of } g$$

then g is the *inverse* of f and we denote g by f^{-1} .

