

MATH 510, FTC

Modern Analysis

James Madison University

Some basic facts about integrals:

1. If $f_1(t)$ and $f_2(t)$ are integrable functions, then

$$\int_a^b f_1(t) + f_2(t) dt = \int_a^b f_1(t) dt + \int_a^b f_2(t) dt$$

2. If f is integrable and a, b, c are three points, then

$$\int_a^b f(t) dt = \int_a^c f(t) dt + \int_c^b f(t) dt$$

3. If f is integrable then so is $|f|$ and

$$\left| \int_a^b f(t) dt \right| \leq \int_a^b |f(t)| dt$$

Theorem If g is a continuous function and G is defined by

$$G(x) = \int_a^x g(t) dt$$

the G is differentiable and

$$G'(x) = g(x).$$

proof We need to consider

$$\lim_{\Delta x \rightarrow 0} \frac{G(x + \Delta x) - G(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\int_a^{x+\Delta x} g(t) dt - \int_a^x g(t) dt}{\Delta x}$$

and show that this limit is $g(x)$. Applying “basic fact” number 2 above with $c = x$ and $b = x + \Delta x$ we have

$$\lim_{\Delta x \rightarrow 0} \frac{G(x + \Delta x) - G(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\int_x^{x+\Delta x} g(t) dt}{\Delta x}$$

So then, we need to show that given any $\epsilon > 0$ we can find a corresponding $\delta > 0$ so that $|g(x) - \frac{\int_x^{x+\Delta x} g(t) dt}{\Delta x}| < \epsilon$ when $|\Delta x| < \delta$. But we know that since g is continuous, given $\epsilon > 0$ we can find a corresponding $\delta > 0$ so that $|g(t) - g(x)| < \epsilon$ when $|x - t| < \delta$. Thus:

$$\begin{aligned}
\left| g(x) - \frac{\int_x^{x+\Delta x} g(t) dt}{\Delta x} \right| &= \left| g(x) - \frac{\int_x^{x+\Delta x} g(t) - g(x) + g(x) dt}{\Delta x} \right| \\
&= \left| g(x) - \frac{\int_x^{x+\Delta x} g(t) - g(x) dt + g(x) \int_x^{x+\Delta x} dt}{\Delta x} \right| \\
&= \left| g(x) - \frac{\int_x^{x+\Delta x} g(t) - g(x) dt + g(x) \cdot \Delta x}{\Delta x} \right| \\
&= \left| \frac{\int_x^{x+\Delta x} g(t) - g(x) dt}{\Delta x} \right| \\
&\leq \frac{\int_x^{x+\Delta x} |g(t) - g(x)| dt}{\Delta x} < \frac{\int_x^{x+\Delta x} \epsilon dt}{\Delta x} = \epsilon
\end{aligned}$$

Since g is continuous, given $\epsilon > 0$ we can find a corresponding $\delta > 0$ so that $|g(t) - g(x)| < \epsilon$ when $|x - t| < \delta$.