Statistical **inference** draws conclusions about a population from sample data.

Conditions about estimating a population mean μ :

- 1. We have a SRS from the population.
- 2. The variable has a normal distribution $N(\mu, \sigma)$ or n > 30.

3. We know population standard deviation σ (this assumption is unrealistic and will be relaxed later).

< 口 > < 同 > < 回 > < 回 > < 回 > <

A **point estimate** is a single number that is used to estimate an unknown parameter.

A level $(1 - \alpha)100\%$ confidence interval for a parameter has 1. an interval of the form point estimate \pm margin of error 2. a confidence level $(1 - \alpha)100\%$ which gives the success rate for the method.

< 口 > < 同 > < 回 > < 回 > < 回 > <

Confidence interval

$$\bar{x} \pm z \frac{\sigma}{\sqrt{n}}$$

where $z = z_{\alpha/2}$. Common critical values: 90% : $z_{0.05} = 1.645$ 95% : $z_{0.025} = 1.96$ 99% : $z_{0.005} = 2.576$ Example 7.3.

The sampling distribution of \bar{x}



Note here $\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$ The empirical rule says 95% of the \bar{x} s are within 2 (1.96 to be exact) $\sigma_{\bar{x}}$ of μ , so a margin of error $2\sigma_{\bar{x}}$ can guarantee a 95% success rate.

Find the z critical value



(James Madison University)

October 16, 2020 5/47

э

イロト イヨト イヨト イヨト



use $p=\frac{1-ConfidenceLevel}{2}$ or $p=\frac{1+ConfidenceLevel}{2}$ in the z chart to find z. e.g., for 92% confidence interval, use $p=\frac{1+0.92}{2}=0.96$ to find z=1.75. for 88% confidence interval, use $p=\frac{1+0.88}{2}=0.94$ to find z=1.55.

Example 7.5. A simple random sample of 6 cereal boxes produces $\bar{x} = 20.25$ ounces. The fill weights are normally distributed with $\sigma = 0.2$. Get a 90% CI for the mean fill weight. $z_{0.05} = 1.645$, the CI is $\bar{x} \pm 1.645 \frac{\sigma}{\sqrt{n}} = 20.25 \pm 1.645 \frac{0.2}{\sqrt{6}} = 20.25 \pm 0.1343 = (20.12, 20.38)$ oz.

イロト イ団ト イヨト イヨト

Exercises

A random sample of n = 360 three month boys produce a sample mean weight $\bar{x} = 25.5$ pounds. Suppose the population standard deviation is given as $\sigma = 5.3$.

Get a 95% CI for the population mean weight.

< 口 > < 同 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ >

Solutions

$$z_{0.025} =$$
 1.96,
25.5 \pm 1.96 $*$ $rac{5.3}{\sqrt{360}} =$ 25.5 \pm 0.55 $=$ (24.95, 26.05) lb.

(James Madison University)

2

イロト イヨト イヨト イヨト

Section 7.1 prob. 43.

The mean math SAT score of a random sample of 100 freshmen is 458. Assume $\sigma = 116$. Find a 99% CI for the population mean.

 $458 \pm 2.576 \frac{116}{\sqrt{100}} = 458 \pm 29.88 = (458.12, 487.88).$ For confidence level=0.99, use p = 0.995 in the z chart to find z = 2.57 or z = 2.58, a more exact value is z = 2.576.

Margin of error

Note margin of error (denoted by m) in the confidence interval is

$$m = z \frac{\sigma}{\sqrt{n}}$$

sample size needed to achieve a certain margin of error:

$$n = (\frac{z\sigma}{m})^2$$

< ロ > < 同 > < 回 > < 回 >

The weight of mice fed a certain diet is normally distributed with $\sigma = 3$ grams. How many mice must be weighted so than a 95% CI will have a margin of error of 0.5 grams? $0.5 = 1.96\frac{3}{\sqrt{2}}$

or

$$n = (\frac{1.96 * 3}{0.5})^2 = 138.30 \approx 139$$

< 口 > < 同 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ >

October 16, 2020

12/47

We always round up *n* when computing the needed sample size.

Exercise 51

In a random sample of 100 electronic components, the mean life time was 125 hours. Assume that the components lifetimes are normally distributed with population standard deviation $\sigma = 20$ hours.

a). Construct a 98% CI for the mean battery life.

b). Find the sample size needed so that a 99% CI will have a margin of error of 3.

・ ロ ト ・ 同 ト ・ 回 ト ・ 回 ト

Solutions

a).
$$z_{0.01} = 2.33, \bar{x} \pm z \frac{\sigma}{\sqrt{n}}$$

= $125 \pm 2.33 \frac{20}{\sqrt{100}} = 125 \pm 4.66 = (120.34, 129.66).$
b) $z_{0.005} = 2.576, n = (\frac{z\sigma}{m})^2 = (\frac{2.576*20}{3})^2 = 295.$

(James Madison University)

October 16, 2020 14/47

2

イロト イヨト イヨト イヨト

Unknown population standard deviation

Conditions: SRS Population distribution is normal or n > 30. We do not know σ

・ロト ・ 四ト ・ ヨト ・ ヨト

Replace σ with s

Recall
$$\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$$
.
 $z = \frac{\bar{x}-\mu}{\sigma/\sqrt{n}} \sim Normal(0, 1)$.
Now $t = \frac{\bar{x}-\mu}{s/\sqrt{n}}$ has a t distribution with $n - 1$ degrees of freedom.

(日)

October 16, 2020

16/47

(James Madison University)

t distribution



(James Madison University)

э

t and z critical value



level $(1 - \alpha)100\%$ confidence interval: $\bar{x} \pm t_{\alpha/2} \frac{s}{\sqrt{n}}$. Example 7.12. A sample of 123 people reported the number of hours spent on internet per week, $\bar{x} = 8.20, s = 9.84$ hours. Get a 95% CI for μ . $\bar{x} \pm t_{\alpha/2} \frac{s}{\sqrt{n}} = 8.20 \pm 1.984 \frac{9.84}{\sqrt{123}} = 8.20 \pm 1.7603 = (6.44, 9.96)$ hours. Exercise 29, 31.

・ ロ ト ・ 同 ト ・ 回 ト ・ 回 ト

Exercise

The healing rate of wounds made on 18 newts was measured with a sample mean of 25.67 micrometers per hour and a standard deviation s = 8.324. A stem-and-leaf plot of data does not show obvious departure form normality. Get a 95% CI for the population mean.

< 口 > < 同 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ >

t Table

cum. prob	t.50	t.75	t.80	t.as	t_90	t.95	t.975	t.20	t.995	t.999	t_2225
one-tail	0.50	0.25	0.20	0.15	0.10	0.05	0.025	0.01	0.005	0.001	0.0005
two-tails	1.00	0.50	0.40	0.30	0.20	0.10	0.05	0.02	0.01	0.002	0.001
df											
1	0.000	1.000	1.376	1.963	3.078	6.314	12.71	31.82	63.66	318.31	636.62
2	0.000	0.816	1.061	1.386	1.886	2.920	4.303	6.965	9.925	22.327	31.599
3	0.000	0.765	0.978	1.250	1.638	2.353	3.182	4.541	5.841	10.215	12.924
4	0.000	0.741	0.941	1.190	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	0.000	0.727	0.920	1.156	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	0.000	0.718	0.906	1.134	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	0.000	0.711	0.896	1.119	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	0.000	0.706	0.889	1.108	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	0.000	0.703	0.883	1.100	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	0.000	0.700	0.879	1.093	1.372	1.812	2.228	2.764	3.169	4.144	4.587
11	0.000	0.697	0.876	1.088	1.363	1.796	2.201	2.718	3.106	4.025	4.437
12	0.000	0.695	0.873	1.083	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	0.000	0.694	0.870	1.079	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	0.000	0.692	0.868	1.076	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	0.000	0.691	0.866	1.074	1.341	1.753	2.131	2.602	2.947	3.733	4.073
16	0.000	0.690	0.865	1.071	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	0.000	0.689	0.863	1.069	1.333	1.740	2.110	2.567	2.898	3.646	3.965
18	0.000	0.688	0.862	1.067	1.330	1.734	2.101	2.552	2.878	3.610	3.922
19	0.000	0.688	0.861	1.066	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	0.000	0.687	0.860	1.064	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	0.000	0.686	0.859	1.063	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	0.000	0.686	0.858	1.061	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	0.000	0.685	0.858	1.060	1.319	1.714	2.069	2.500	2.807	3.485	3.768
24	0.000	0.685	0.857	1.059	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	0.000	0.684	0.856	1.058	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	0.000	0.684	0.856	1.058	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	0.000	0.684	0.855	1.057	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	0.000	0.683	0.855	1.056	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	0.000	0.683	0.854	1.055	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	0.000	0.683	0.854	1.055	1.310	1.697	2.042	2.457	2.750	3.385	3.646
40	0.000	0.681	0.851	1.050	1.303	1.684	2.021	2.423	2.704	3.307	3.551
60	0.000	0.679	0.848	1.045	1.296	1.671	2.000	2.390	2.660	3.232	3.460
80	0.000	0.678	0.846	1.043	1.292	1.664	1.990	2.374	2.639	3.195	3.416
100	U.000	0.677	0.845	1.042	1.290	1.660	1.984	2.364	2.626	3.174	3.390
1000	0.000	0.675	0.842	1.037	1.282	1.646	1.962	2.330	2.581	3.098	3.300
z	0.000	0.674	0.842	1.036	1.282	1.645	1.960	2.326	2.576	3.090	3.291
L	0%	50%	60%	70%	80%	90%	95%	98%	99%	99.8%	99.9%
	Confidence Level										

◆□> ◆圖> ◆理> ◆理> 「理

Table A.2 Cumulative Normal Distribution (continued)



z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998
3.5	.9998	.9998	.9998	.9998	.9998	.9998	.9998	.9998	.9998	.9998
3.6	.9998	8999.	.99999	.99999	.99999	.99999	.99999	.99990	.9999	9999

(James Madison University)

◆□> ◆□> ◆注> ◆注> □注

October 16, 2020

22/47

Solution

n=18, d.f. = 17, 25.67
$$\pm$$
 2.110 $\frac{8.324}{\sqrt{18}}$ = 25.67 \pm 4.14 = (21.53, 29.81) hours.

<ロ> (日) (日) (日) (日) (日)

2

23/47

October 16, 2020

(James Madison University)

Gas bubbles inside ancient amber gives the following nitrogen (percent). 63.4, 65.0, 64.4,63.3, 54.8, 64.5, 60.8, 49.1, 51.

 $\bar{x} = 59.59, s = 6.26.$

Get a 90% CI for μ , the mean nitrogen content in ancient air.

< 口 > < 同 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ >

October 16, 2020

24/47

CI: $59.59 \pm 1.860 * \frac{6.26}{\sqrt{9}} = 59.59 \pm 3.88 = (55.71, 63.47).$

12

Six measurements were made of the mineral content (in percent) of spinach. It is reasonable to assume the population is approximately normal:

19.1, 20.8, 20.8, 21.4, 20.5, 17.4.

Get a 95% CI for the mean mineral content.

 $\bar{x} = 20.0, s = 1.49,$ CI : $20.0 \pm 2.571 * 1.49/\sqrt{6} = 20 \pm 1.56 = (18.44, 21.56).$

Note: $20.0 \pm 2.571 * 1.49 / \sqrt{6} = 20.0 \pm (2.571 * 1.49 / \sqrt{6}).$

< 日 > < 同 > < 回 > < 回 > < □ > <

One sample t test

 $H_0: \mu = \mu_0.$ test statistic $t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}.$ example 16.3. exercise 16.8, 16.9, 16.10. 16.10: Amber gas data: test $H_0: \mu = 78.1$ $H_a: \mu \neq 78.1.$

Robustness of t procedures

The conditions for using the t procedures:

- 1. The sample is SRS.
- 2. n < 15: data appear close to normal.
- 3. $n \ge 15$: no outliers or strong skewness.
- 4. $n \ge 40$: okay for skewed distribution.

A B F A B F

Seven subjects who identified themselves as Buddhist reported hours per week watching TV of:

2,1,1,3,2,3,2.

Assume the TV watching time of a budddist is approximately normally distributed.

1). Estimate the sample mean \bar{x} and sample standard deviation *s*.

2). Construct a 90% confidence interval for μ , the mean TV watching time per week for all Buddhists.

 $ar{x} = 2, s = 0.816.$ 2 \pm 1.943 * 0.816/ $\sqrt{7} = (1.40, 2.60)$ hours.

A random sample of emissions is taken from an incinerator for 4 days this year. The observations were:

2, 3, 3, and 2 grams of dioxin per day.

Construct a 90% confidence interval for the mean emission of dioxin per day.

$ar{x} \pm t^* rac{s}{\sqrt{n}} = 2.5 \pm 2.353 * 0.577/\sqrt{4} = (1.82, 3.18)$ grams.

<ロ> <四> <四> <四> <四> <四</p>

October 16, 2020

32/47

(James Madison University)

sampling distribution of \hat{p}

population proportion: p sample proportion: $\hat{p} = \frac{x}{n}$ x : number of successes in the sample. $\mu_{\hat{p}} = p, \sigma_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}.$

For large *n*, approximately $\hat{p} \sim \text{Normal } (p, \sqrt{\frac{p(1-p)}{n}}).$

Sample size needed: $n\hat{p} \ge 10$ and $n(1 - \hat{p}) \ge 10$. (at least 10 successes and 10 failures in the sample) Data come from a random sample.

$$(1 - \alpha)$$
100%Cl is:
 $\hat{p} \pm z \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}.$

3

< 日 > < 同 > < 回 > < 回 > < □ > <

When the 2000 GSS asked subjects whether they would be willing to accept cuts in their standard of living to protect the environment, 344 of 1170 subjects said yes. Find a 95% confidence interval for the population proportion who would say yes.

The sample proportion is $\hat{p} = 344/1170 = 0.294$ A 95% CI:

$$\hat{p} \pm 1.96 * \sqrt{rac{\hat{p}(1-\hat{p})}{n}} = 0.294 \pm 1.96 \sqrt{rac{0.294 * 0.706}{1170}} = 0.294 \pm 0.026 = (0.268, 0.320).$$
exercise: 23, page 327.

< 口 > < 同 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ >



In 2008 a GSS asked 182 people whether they received health insurance from their employers. A total of 60 people said they did. Get a 95% CI for the population proportion.

< ロ > < 同 > < 回 > < 回 >

Solution

$$\hat{p} = 60/182 = 0.330.$$

95% CI: $0.330 \pm 1.96 \sqrt{\frac{0.330 * 0.670}{182}} = 0.330 \pm 0.068 = (0.262, 0.398).$

2

イロト イヨト イヨト イヨト

sample size for desired margin of error

$$m = z \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$
, and

$$n = \hat{p}(1-\hat{p})\big(\frac{z}{m}\big)^2$$

< 口 > < 同 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ >

October 16, 2020

38/47

Use $\hat{p} = 1/2$ if a value for \hat{p} is unavailable.

(James Madison University)

Example 7.14

Among a sample of 517 music teachers, 403 believe that video games have a positive effect on music education. Estimate the sample size needed so that a 95% CI will have a margin of error of 0.03.

$$\hat{p} = rac{403}{517} = 0.7795,$$

 $n = \hat{p}(1-\hat{p})(rac{z}{m})^2 = 0.7795 * (1-0.7795)(rac{1.96}{0.03})^2 \approx 734.$

exercise 30 page 328:

1). A gallup poll estimates the proportion of people who believe the economy is getting better to be 0.33. What sample size is needed so that the 95% CI will have a margin of error 0.03?

イロト 不得 トイヨト イヨト

October 16, 2020

39/47

2).Estimate sample size needed if no estimate of p is available.

Solutions

a.
$$n = 0.33 * 0.67 (\frac{1.96}{0.03})^2 = 944$$
.

b.
$$n = 0.50 * 0.50(\frac{1.96}{0.03})^2 = 1068$$
.

イロト イヨト イヨト イヨト

2

40/47

October 16, 2020

(James Madison University)



A researcher wants to estimate the population proportion of frogs with fungus infection in a pond. Her sample proportion is \hat{p} =0.35 based on a sample of 100.

- 1). Construct a 95% confidence interval for the population proportion *p*.
- 2). Construct a 99% confidence interval for *p*.

Solutions

1).
$$0.35 \pm 1.96 \sqrt{\frac{0.35 \pm 0.65}{100}} = 0.35 \pm 0.09 = (0.26, 0.44).$$

2). $0.35 \pm 2.576 \sqrt{\frac{0.35 \pm 0.65}{100}} = 0.35 \pm 0.12 = (0.23, 0.47).$

イロト イヨト イヨト イヨト

2

42/47

October 16, 2020

(James Madison University)

adjusted sample proportion: $\tilde{p} = \frac{x+2}{n+4}$ Replace \hat{p} and n with \tilde{p} and $\tilde{n} = n+4$ in the CI formula. $\tilde{p} \pm z \sqrt{\frac{\tilde{p}(1-\tilde{p})}{\tilde{n}}}$



2. A 2008 GSS survey asked 182 whether they received health insurance from their employers and 60 of them said yes. Get a 90% CI for the population proportion who receive health insurance from their employers.

< 口 > < 同 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ >

$\hat{p} = 60/182 = 0.330.$ 90% CI: $0.330 \pm 1.645 \sqrt{\frac{0.330 * 0.670}{182}} = 0.330 \pm 0.057 = (0.273, 0.387).$

< 🗇 🕨



The General Social Survey recently asked 1294 people whether they performed any volunteer work during the past year. A total of 517 people said they did.

1). Find a point estimate for the proportion of people who performed volunteer work during the past year.

2). Construct a 95% confidence interval for the proportion of people who performed volunteer work during the past year.

3). A sociologist states that 50% of Americans perform volunteer work in a given year. Does the confidence interval contradict his statement? Explain.

October 16, 2020

46/47

Solutions

1). $\hat{p} = \frac{517}{1294} = 0.400$. 2). $0.400 \pm 1.96\sqrt{.400 * .600/1294} = 0.400 \pm 0.027 = (0.373, 0.427)$. 3). Yes, the confidence interval contradicts his statement as .50 is not included in the interval.

・ロト ・ 四ト ・ ヨト ・ ヨト …